

# Update on Canadian Guidelines on the Prevention, Assessment and Treatment of Depression among Older Adults

Dr. David Conn

# Objectives

- **By the end of the presentation participants will:**
  1. Be aware of the process for creating and updating the Guidelines
  2. Be able to consider possible approaches to the prevention of depression in older adults.
  3. Consider utilizing updated recommendations regarding the treatment of depression in their practice.

# Disclosures

- No disclosures related to this presentation
- Funding to CCSMH for the original guidelines was provided by Public Health Agency of Canada.
- Guideline update funding was provided by a CIHR-Institute of Aging Betty Havens Award for KT in Aging.

# Global prevalence and burden of depressive and anxiety disorders in 204 countries and territories in 2020 due to the COVID-19 pandemic



COVID-19 Mental Disorders Collaborators\*



www.thelancet.com Published online October 8, 2021 [https://doi.org/10.1016/S0140-6736\(21\)02143-7](https://doi.org/10.1016/S0140-6736(21)02143-7)





Figure 2: Change in the prevalence of major depressive disorder after adjustment for (ie, during) the COVID-19 pandemic, 2020



**Figure 4: Global burden of major depressive disorder and anxiety disorders by age and sex, 2020**  
 Baseline refers to pre-pandemic DALYs and additional refers to additional burden due to the COVID-19 pandemic.  
 DALYs=disability-adjusted life-years.



## Canadian Coalition for Seniors' Mental Health (CCSMH)


[www.ccsmh.ca](http://www.ccsmh.ca)

**The mission of the CCSMH is:**

To promote the mental health of older adults by connecting people, ideas, and resources.


Founded 2002 - Linked to CAGP

# The Bio-Psycho-Social Model



## Canadian Guidelines on Prevention, Assessment and Treatment of Depression Among Older Adults

ccsmh.ca



## 2 main sections:

- New or modified recommendations
- Side-by-side comparison of 2006 and 2021 recommendations (50 / 71 essentially unchanged)
- **Ideally readers will review both sections**

# Topics included

1. Prevention
2. Assessment / Screening
3. Psychosocial interventions and Psychotherapy
4. Pharmacological treatments
5. Somatic Treatments
6. Subtypes of depression
7. Special populations
8. Models of Care

## 2021 Guideline Update Group

**Dr. David Conn, M.B., B.Ch., B.A.O.,  
FRCPC**

***Group Chair***

VP, Education and Director of Centre  
for Education, Baycrest Health  
Sciences  
Professor, Department of Psychiatry,  
University of Toronto  
Co-Chair, Canadian Coalition for  
Seniors' Mental Health

**Lori Amdam, B.Sc.N, M.S.N.**

***Group Member***

Education Consultant, Gerontology  
CCSMH Steering Committee  
Member  
Vancouver Island, British Columbia,  
Canada

**Dr. Daniel M. Blumberger, MD, MSc,  
FRCPC**

***Group Member***

Director & Chair, Temerty Centre for  
Therapeutic Brain Intervention  
Clinician Scientist, Campbell Family  
Research Institute  
Centre for Addiction and Mental  
Health  
Professor, Department of Psychiatry,  
University of Toronto

**Dr. Chris Frank, MD, FCFP**

***Group Member***

Care of the Elderly Family Physician,  
Professor, Department of Medicine,  
Queen's University, Kingston,  
Ontario

**Dr. Cindy Grief, MSc., MD, FRCPC**  
***Group Member***

Medical Director, Mental Health  
Services, Baycrest Health Sciences  
Assistant Professor,  
University of Toronto

**Dr. Robert Madan, MD, FRCPC**

***Group Member***

Psychiatrist-in-Chief & Executive  
Medical Director, Baycrest Health  
Sciences  
Associate Professor,  
University of Toronto

**Dr. Kiran Rabheru, MD, CCFP, FRCPC**

***Group Member***

Geriatric Psychiatrist, TOH  
Professor of Psychiatry,  
University of Ottawa  
Co-Chair, Canadian Coalition for  
Seniors' Mental Health

**Anar Dilara MBBS, MPH, MPhil**

Former Research Assistant,  
Canadian Coalition for Seniors'  
Mental Health

**Daniella Minchopoulos, BSc**

Former Project Assistant,  
Canadian Coalition for Seniors'  
Mental Health

## Original 2006 Guideline Development Group (with 2006 titles)

**Dr. Marie-France Tourigny-Rivard, M.D., FRCPC**  
*Co-Lead*  
Professor and Chief of the Geriatric Psychiatry Division,  
Department of Psychiatry,  
University of Ottawa  
Clinical Director, Geriatric Psychiatry Program, Royal Ottawa Hospital, Ottawa, Ontario

**Dr. Diane Buchanan, RN, DNSc, GNC (C)**  
*Co-Lead*  
Assistant Professor, School of Nursing, Queen's University, Kingston, Ontario

**Dr. Philippe Cappeliez, Ph.D.**  
*Group Member*  
Professor, School of Psychology,  
University of Ottawa,  
Ottawa, Ontario

**Dr. Chris Frank, MD, FCFP**  
*Group Member*  
Care of the Elderly, Professor,  
Department of Medicine,  
Queen's University,  
Kingston, Ontario

**Pronica Janikowski, R.Ph., B.Sc. Phm., CGP**  
*Group Member*  
Certified Geriatric Pharmacist,  
Long-Term Care Facility Consultant,  
Picton Clinic Pharmacy,  
Picton, Ontario

**Faith Malach, MHSc, MSW, RSW**  
*Project Director*  
Executive Director, Canadian Coalition for Seniors' Mental Health,  
Adjunct Practice Professor, Faculty of Social Work, University of Toronto, Toronto, Ontario

**Jennifer Mokry, MSW, RSW**  
*Project Coordinator*  
Project Manager, Canadian Coalition for Seniors' Mental Health,  
Toronto, Ontario

**Lily Spanjevic, RN, BScN, MN, CPMHN(C), GNC (C)**  
*Group Member*  
Advanced Practice Leader-Geriatric Rehabilitation Program,  
Toronto Rehabilitation Institute,  
Toronto, Ontario

**Dr. Alastair Flint, MB, FRCPC, FRANZCP**  
*Consultant*  
Professor of Psychiatry, University of Toronto  
Head, Geriatric Psychiatry Program, University Health Network, Toronto, Ontario

**Dr. Nathan Herrmann, MD, FRCPC**  
*Consultant*  
Head, Division of Geriatric Psychiatry, Sunnybrook and Women's College Health Sciences Centre, Professor of Psychiatry, University of Toronto, Toronto, Ontario

**A** systematic search for peer-reviewed scholarly articles was performed in 5 databases including Medline, Embase, HealthStar, Cochrane, and PsychINFO from the beginning of the end date of the literature review from the previous version of the guideline—July 2006—through December 2018. The searches included and were restricted to English papers only and search terms used were the same as for the 2006 Guidelines.

Multiple phases of title and abstract review were conducted by one of the authors to identify 344 full-text articles from an initial yield of 1560 articles from database searches; those were further categorized based on types of study such as controlled trials (especially randomized), meta-analyses, reviews (especially systematic), and practice guidelines or expert committee reports potentially relevant to the subject area. Subsequently, additional relevant articles of which the members were aware were included.

## CATEGORIES OF EVIDENCE FOR CAUSAL RELATIONSHIPS AND TREATMENT

Evidence from meta-analysis of randomized controlled trials	Ia
Evidence from at least 1 randomized controlled trial	Ib
Evidence from at least 1 controlled study without randomization	IIa
Evidence from at least 1 other type of quasi-experimental study	IIb
Evidence from non-experimental descriptive studies, such as comparative studies, correlation studies, and case-control studies	III
Evidence from expert committees reports or opinions and/or clinical experience of respected authorities	IV

(Shekelle et al., 1999)

## STRENGTH OF RECOMMENDATION

Directly based on category I evidence	A
Directly based on category II evidence or extrapolated recommendation from category I evidence	B
Directly based on category III evidence or extrapolated recommendation from category I or II evidence	C
Directly based on category IV evidence or extrapolated recommendation from category I, II, or III evidence	D

(Shekelle et al., 1999)

# Differential Diagnosis “I feel down...”

- Major Depressive Disorder – Major Depressive Episode
- Bipolar Disorder – Major Depressive Episode
- Persistent Depressive Disorder- pure dysthymic type
- “Minor/subthreshold depression”
- Adjustment Disorder
- Bereavement
- Personality Disorder
- Mood disorder due to a medical condition
- Substances/meds

# MDE Symptoms (DSM-5)

- \*Depressed/empty
- \*Anhedonia
- Sleep (less or more)
- Appetite (less or more)
- Reduced energy
- Poor concentration
- Guilt or worthlessness
- Psychomotor slowing or restlessness
- Thoughts of death or suicide
- Distress or impairment in functioning X 2 weeks
- 5/9 symptoms

Depressed For  
No Reason



Depressed For  
A Good Reason




JUST  
DEPRESSED,  
DON'T  
WANT TO  
ANALYZE  
IT →

Still Depressed

UN-  
DONE  
→

# Recommendations: Prevention





*An ounce of prevention  
is worth a pound of cure*  
- Ben Franklin

## What Makes Us Healthy



## What We Spend On Being Healthy



Data courtesy of the Boston Foundation and the New England Healthcare Institute

# Prevention

- Although research on prevention is in its infancy, prevention may be an alternative strategy to further reduce the disease burden of depression, which has been described as a global public health priority (Reynolds et al., 2017 ).

*Universal* prevention focuses on the general public or a whole population group regardless of risk status.

*Selective* prevention targets individuals or subgroups that are at higher risk of developing mental disorders than average individuals or subgroups.

*Indicated* prevention focuses on individuals who are identified as having prodromal symptoms or biological markers of mental disorders, but who do not yet meet the diagnostic criteria for a full-blown diagnosis.




Fig. 1. Strength of evidence on risk factors for cognitive decline.

# Potentially Modifiable Risk Factors for Dementia

1. Education
2. Hearing Loss
3. Obesity
4. Hypertension
5. Smoking
6. Physical inactivity
7. Diabetes
8. Depression
9. Social isolation

- Interventions might have the potential to delay or prevent a third of dementia cases”

**NEW:**

## **PREVENTION**

**A variety of interventions focused on reducing social isolation and/or loneliness in older adults have demonstrated a reduction in depressive symptoms in addition to reduced loneliness. These interventions were primarily group-based and in long-term care settings. They include reminiscence therapy, physical exercise programs, videoconferences with family, horticultural therapy, and gender-based social groups. [B]**

Systematic reviews: Quan et al., (2019), *Aging Mental Health*,  
Franck et al., (2016), *Qual Life Res*.

**NEW:**

## **PREVENTION**

**Social prescribing, which is defined as, “a means of enabling primary care services to refer patients with social, emotional, or practical needs to a range of local, non-clinical services, often provided by the voluntary and community sector”, may result in reduced depressive symptoms among older adults who have experienced mild to moderate symptoms of depression, social isolation, or loneliness. [C]**

**NEW:**

**PREVENTION**

**A stepped-care approach (e.g., watchful waiting, cognitive behavioural therapy [CBT-based] bibliotherapy, problem-solving therapy, and referral to primary care for antidepressant medication) can reduce the incidence of depressive and anxiety disorders in community-dwelling older adults with subthreshold depression or anxiety. [B]**

Van't Veer-Tazelaar et al. (2009) studied the following steps: a watchful waiting approach, CBT-based bibliotherapy, CBT-based problem-solving treatment, and referral to primary care for medication, if required. **The intervention group had a 50% reduction in incidence of major depressive disorder or anxiety disorder over a 12-month period compared to usual care.**

**NEW:**

## **PREVENTION**

**Higher levels of physical activity are consistently associated with lower odds of developing future depression. This finding is consistent across all age groups including older adults. Clinicians should encourage patients with low levels of physical activity to become more active. Tools are available for clinicians to assist patients in setting health-related goals (e.g., Fountain of Health). [B]**

Schuch et al. (2018): meta-analysis of 49 studies of physical activity. Compared with people with low levels of physical activity, those with high levels had significantly lower odds of developing depression. **Physical activity had a protective effect against the emergence of depression across all age groups including older adults.**

# The Fountain of Health Initiative

Bringing Seniors' Mental Health Promotion into  
Clinical Practice



**fountain** *of* **health**

# FoH 5 Key Areas of Focus



fountain of health

Social Activity

Positive Thinking

Physical Activity

Mental Health

Lifelong Learning

**Clinicians should utilize the instilling of hope and positive thinking as important therapeutic tools in the prevention of depression and in helping individuals with depressive symptoms or disorders. [D]**

A review of 9 studies, utilizing such interventions by nurses in patients coping with cancer, concluded that it is possible to increase hope in this group (Li et al., 2018). Moore (2005) suggests that nurses are in key positions to have conversations with their patients about hope and about strategies to find renewed hope in any situation. **Although more research is necessary to understand optimal interventions, we would encourage all healthcare staff to reflect on how best to incorporate the instilling of hope into their practices.**

# Steps to develop a positive attitude

- Look at the bright side of life
- Have faith in yourself and the power of the Universe
- Associate with positive people
- Read inspiring stories
- Repeat affirmations that inspire and motivate
- Visualize only what you want to happen
- Learn to master your thoughts
- Meditate

R. Sasson

# SCREENING & ASSESSMENT

## Recommendations: Screening and Assessment – Risk Factors

2006 RECOMMENDATIONS	2021 NEW OR UPDATED RECOMMENDATIONS
<p>Health care providers should be familiar with the physical, psychological, and social risk factors for depressive disorders in older adults and include a screening for depression for their clients/ patients who present with some of these risk factors. <b>[D]</b></p>	<p>Unchanged</p>
<p>We recommend targeted screening of those elderly at higher risk for depression due to the following situations:</p> <ul style="list-style-type: none"><li>• Recently bereaved with unusual symptoms (e.g., active suicidal ideation, guilt not related to the deceased, psychomotor retardation, mood congruent delusions, marked functional impairment after 2 months of the loss, reaction that seems out of proportion with the loss)</li><li>• Bereaved individuals, 3 to 6 months after the loss</li><li>• Socially isolated</li><li>• Persistent complaints of memory difficulties</li><li>• Chronic disabling illness</li><li>• Recent major physical illness (e.g., within 3 months)</li><li>• Persistent sleep difficulties</li><li>• Significant somatic concerns or recent onset anxiety</li><li>• Refusal to eat or neglect of personal care</li><li>• Recurrent or prolonged hospitalization</li><li>• Diagnosis of dementia, Parkinson disease or stroke</li><li>• Recent placement in a nursing/Long Term Care (LTC) home <b>[B]</b></li></ul>	<p>Essentially unchanged, but note that the exclusion of major depression in the first 2 months after the loss and in bereavement was removed in Diagnostic and Statistical Manual of Mental Disorders, 5<sup>th</sup> Edition (DSM-V).</p>

**Recommendations: Screening and Assessment – Screening and Screening Tools**

2006 RECOMMENDATIONS	2021 NEW OR UPDATED RECOMMENDATIONS
Appropriate depression screening tools for elderly persons without significant cognitive impairment in general medical or geriatric settings include the self-rating <i>Geriatric Depression Scale (GDS)</i> , the <i>SELFCARE self-rating scale</i> , and the <i>Brief Assessment Schedule Depression Cards (BASDEC)</i> for hospitalized patients. <b>[B]</b>	We recommend using the GDS or the Patient Health Questionnaire-9 (PHQ-9). <b>[B]</b>
For patients with moderate to severe cognitive impairment, an observer-rated instrument, such as the Cornell Scale for Depression in Dementia is recommended instead of the GDS. <b>[B]</b>	Unchanged

# Assessment

- Interview
- Collateral History
- Safety
- Cognitive Assessment
- Physical exam
- Bloodwork
  - CBC, Lytes, Thyroid, B12, Calcium, glucose
- Imaging
- Driving

Recommendations:  
Psychotherapy and  
psychosocial interventions



**NEW:**

## **PSYCHOTHERAPIES AND PSYCHOSOCIAL INTERVENTIONS**

There is promising evidence for exercise and mind-body interventions (e.g., tai chi, yoga, and mindfulness-based stress reduction) in reducing depressive symptoms in late-life either alone or in combination with other therapies. Physical activity in the form of exercise is an important non-pharmacological approach to improve mood in older adults. Clinicians should use their judgement in recommending the type of exercise and duration, taking into account comorbidities, physical capacity, and level of motivation. **[B]**

**MODIFIED:****PSYCHOTHERAPIES AND PSYCHOSOCIAL INTERVENTIONS**

Psychotherapies with the most evidence for effectiveness in older adults include: cognitive behaviour therapies (CBT; individual and group) and problem-solving therapy (PST). PST can be provided to older adults with cognitive impairment and executive dysfunction; CBT and PST have also shown benefit for older adults with depression and medical comorbidity. [B]

Psychotherapies and psychosocial treatments should be made available to older adults with depression (symptoms and disorder) in diverse settings (community, hospital, long-term care) across all regions of Canada. [A]

There is also evidence to support behaviour therapy, behavioural activation, reminiscence, and other psychotherapies including psychodynamic psychotherapy and interpersonal psychotherapy (IPT). [B]

Internet-delivered therapies may be comparable to face-to-face treatment, and may improve access to services for individuals in under-serviced areas and those with mobility issues. [C]

# Recommendations: Antidepressant medication



**MODIFIED:****SELECTING AN ANTIDEPRESSANT/MONITORING FOR SIDE EFFECTS AND DRUG INTERACTIONS**

It is recommended that clinicians consider sertraline or duloxetine as first-line medications for an acute episode of major depression in older adults. Alternatives include escitalopram and citalopram based on the low possibility of drug interactions but concern about QTc interval may limit dosage to sub-therapeutic levels. [A]

In addition, we suggest clinicians should choose an antidepressant with lowest risk of anticholinergic side effects and drug-drug interactions, as well as being relatively safe in the case of cardiovascular comorbidity. Patients need to be closely monitored for medication compliance, substance use, suicidal ideation, and development of drug toxicity. [D]

**MODIFIED:**

## **TITRATION AND DURATION OF THERAPY (FREQUENCY OF FOLLOW-UP)**

**When starting antidepressants, patients should initially be seen every 1–2 weeks (in-person or virtually) to assess response, side effects, and to titrate the dose. Visits should include, at a minimum, supportive psychosocial interventions and monitoring for worsening of depression, agitation, and suicide risk. [D]**

**MODIFIED:****MONITORING FOR SIDE EFFECTS AND DRUG INTERACTIONS (SODIUM)**

When prescribing SSRI or SNRI antidepressants to older adults, the prescriber should screen for a history of hyponatremia before prescribing, as part of the consent process and then consider getting a sodium level prior to starting the antidepressant if there is a history of hyponatremia. [C]

A serum sodium level should be done within 2–4 weeks of initiating SSRI or SNRI antidepressants. Prescribers may consider checking the level after 2 weeks for those patients on diuretics or who have a history of hyponatremia. There is a lower risk of hyponatremia with TCAs, bupropion, and mirtazapine. [C]

# How Long to Stay on Antidepressant?

- Minimum of 1 year, even in the case of a single episode (Diniz 2014).
- A Cochrane review in 2016
  - quality of evidence was low with only 3 RCTs
  - NNT = 5
  - The authors suggest that “Continuing antidepressant medication for 12 months appears to be helpful with no increased harms..”

# Unchanged

- Older patients who have had more than 2 depressive episodes, had particularly severe or difficult-to-treat depressions or required ECT should continue to take antidepressant maintenance treatment indefinitely, unless there is a specific contraindication to its use.

[D]

# Treatment Resistance

- Only 50% of elderly patients respond to first-line treatment and less than 40% reach remission

# Approach

- Assess medication adherence
- ?alcohol, substances, and medications
- The diagnosis should be reviewed
- Drug-drug interactions
- Medical conditions should be reviewed, hyponatremia.

**MODIFIED:****TITRATION AND DURATION OF THERAPY  
(INADEQUATE RESPONSE)**

When significant improvement has occurred but recovery is not complete after an adequate trial, the clinician should consider:

- a further 4 weeks of monotherapy or consider augmentation with another antidepressant or lithium or an antipsychotic (e.g., aripiprazole) or specific psychotherapy (e.g., IPT, CBT, PST).
- a switch to another antidepressant (same or another class) after discussing with the patient the potential risk of losing any significant improvements made with the first treatment. [C]
- augmentation with lithium remains a viable option but needs to be used carefully due to the risk of lithium toxicity; the clinician must be aware of how to monitor the patient on lithium over time through investigations.

**NOTE:** 2006 recommendation did not include augmentation with an antipsychotic.

# Recommendations regarding subtypes of depression

- Persistent depressive disorder
- MDE – severe without psychosis
- MDE – severe with psychosis

**MODIFIED:****TREATMENT: MAJOR DEPRESSIVE DISORDER,  
SINGLE OR RECURRENT EPISODE – SEVERE BUT  
WITHOUT PSYCHOSIS**

**UNCHANGED** – Patients with severe unipolar depression should be offered a combination of antidepressants and concurrent psychotherapy when appropriate services are available and there is no contraindication to either treatment. [D]

**MODIFIED** – Electroconvulsive therapy (ECT) should be considered in the treatment of older patients with severe unipolar depression who have previously had a good response to a course of ECT and/or failed to respond to 1 or more adequate antidepressant trials plus psychotherapy, especially if their health is deteriorating rapidly due to depression. ECT is a first-line treatment in older, depressed patients who are at high risk of poor outcomes—those with suicidal ideation or intent, severe physical illness, or with psychotic features. [A]

ECT can also be useful for continuation/maintenance therapy of older patients who are partially responsive, treatment resistant, or treatment intolerant with pharmacotherapy during the acute phase of treatment. [B]

**MODIFIED:**

**TREATMENT: MAJOR DEPRESSIVE DISORDER,  
SINGLE OR RECURRENT EPISODES – SEVERE WITH  
PSYCHOTIC FEATURES**

Recently, more placebo-controlled clinical trials reported safe and effective use of combined antidepressant and antipsychotic drugs in MDD with psychotic features, so we recommend that clinicians use their judgement based on severity and patient's physical conditions to try combination pharmacotherapy first. ECT should be considered after 4–8 weeks if combination therapy fails, is poorly tolerated, or if patient develops severe health consequences. [B]



## **NEW: REPETITIVE TRANSCRANIAL MAGNETIC STIMULATION (rTMS)**

**We recommend repetitive transcranial magnetic stimulation (rTMS; left-sided only or sequential bilateral or deep rTMS) for older adults (> 60 years) with unipolar depression who have failed to respond to at least 1 adequate trial of antidepressant. rTMS is not recommended in patients who have failed a course of ECT or who have a seizure disorder. [B]**

**NEW:**

## **PHARMACOGENETIC TESTING**

**At present we do not recommend the broad use of pharmacogenetic testing in older adults with late-life depression. Patients with recurrent severe side effects to several anti-depressant drugs may benefit from pursuing a pharmacogenetic test to see if CYP450 metabolism is contributing. [C]**

# Recommendations: Special Populations-

Depression associated with Dementia,  
Parkinson's Disease & Stroke

# What are the Challenges in Diagnosing Depression in Dementia?

- Unable to recall
- Unable to articulate/describe
- Loss of interest vs inability
- Sleep and appetite
- Psychomotor changes

# Depression of Dementia (dAD) NIMH

## DSM

- At least 5 symptoms
- Almost every day

## dAD

- At least 3 symptoms
- Not every day
- Added irritability
- Added social isolation or withdrawal

**MODIFIED:****SPECIAL POPULATIONS: DEMENTIA**

**UNCHANGED:** Patients who have mild depressive symptoms or symptoms of short duration should be treated with psychosocial supportive interventions first. [D]

**MODIFIED:** There is limited evidence to recommend antidepressant therapy for mild or moderate depression associated with dementia at this time. Behavioural interventions may be utilized as a first-line intervention and antidepressant medication could be offered if symptoms are severe and persistent, understanding that efficacy is not well established and that side effects could occur. [D]

**NEW:**

## **SPECIAL POPULATIONS: PARKINSON'S DISEASE**

**We recommend SSRIs as first line for the treatment of depression in patients with Parkinson's disease with SNRIs as an alternative. CBT can also be considered. [B]**

**MODIFIED:**

**SPECIAL POPULATIONS: VASCULAR DEPRESSION/  
POST-STROKE DEPRESSION**

Consider SSRIs as first-line treatment for post-stroke depression (PSD) regardless of whether or not the stroke is ischemic or hemorrhagic. Second-line treatments can include SNRIs and mirtazapine. Methylphenidate may also be considered, especially if apathy is significant. [B]

**NOTE:** The 2006 recommendations suggested that venlafaxine be avoided. Methylphenidate was not mentioned.



# Recommendations: Models of Care



## MODELS OF CARE

### Recommendations: Models of Care

2006 RECOMMENDATIONS	2021 NEW OR UPDATED RECOMMENDATIONS
<p>Health care professionals and organizations should implement a model of care that addresses the physical/functional as well as the psychosocial needs of older depressed adults.</p> <p>Given the complex care needs of older adults, these are most likely to require interdisciplinary involvement in care, whether in primary care or specialized mental health settings. <b>[B]</b></p>	Unchanged
<p>Health care professionals and organizations should implement a model of care that promotes continuity of care as older adults appear to respond better to consistent primary care providers. <b>[B]</b></p>	Unchanged
	<p><b>New:</b></p> <p>Core elements of evidence-based models for treating late-life depression in primary care include: improved patient education, incorporating interprofessional staff as depression care managers who routinely assess and follow patients clinically, utilizing a stepped-care approach. Treatment prescriptions are provided by a primary care physician or nurse practitioner, with as needed psychiatric consultation. An individualized plan of care should be developed using a collaborative approach. <b>[A]</b></p>
	<p><b>New:</b></p> <p>To optimize access to clinical services, "senior-friendly" virtual care options (e.g., videoconferencing) should be available. Older adult patients should have appropriate equipment and support, to ensure effective and efficient communication to optimize virtual care encounters. <b>[C]</b></p>

# Conclusions

- The updated guidelines have expanded to include prevention strategies
- Greater emphasis on a variety of psychosocial interventions including physical activity
- Some important modifications to pharmacological treatments
- Inclusion of rTMS as a potential treatment
- Updates on evidence-based models of care including virtual care

# Resources



Mental Health  
Commission  
of Canada


Commission de  
la santé mentale  
du Canada



Guidelines for Comprehensive  
Mental Health Services for Older  
Adults in Canada



Canadian Coalition for  
Seniors' Mental Health



Depression in Older Adults:  
a guide for seniors and their families

Depression in Older Adults  
You Are Not Alone!

LEARN MORE + TAKE ACTION

ਵੱਡੀ ਉਮਰ ਦੇ ਲੋਕਾਂ ਵਿੱਚ ਡਿਪ੍ਰੈਸ਼ਨ  
ਤੁਸੀਂ ਇਕੱਲੇ ਨਹੀਂ ਹੋ!

ਹੋਰ ਜਾਣੋ + ਕਾਰਵਾਈ ਕਰੋ

Canadian Coalition for  
Seniors' Mental Health: The  
National Health: The  
depression.





**If you're interested in joining CCSMH  
please contact: [www.ccsmh.ca](http://www.ccsmh.ca)**

**David Conn**

Co-Chair, CCSMH

**Claire Checkland**

Executive Director, CCSMH



**Canadian Coalition for Seniors' Mental Health**

To promote seniors' mental health by connecting people, ideas and resources.

**Coalition Canadienne pour la Santé Mentale des Personnes Âgées**

Promouvoir la santé mentale des personnes âgées en reliant les personnes, les idées et les ressources.

Bring solutions to light  
Mettre en lumière  
les solutions



**Thank You! Any Questions?**

